

Table S1. L-CHOP chemotherapy protocol utilized in this cat during the first course of chemotherapy. This is a 25-week modified University of Wisconsin-Madison chemotherapy protocol.^a

	Week ^b																
	0	1	2	3	4	6	7	8	9	11	13	15	17	19	21	23	25
Vinc^c		x		x		x		x		x		x		x		x	
Cyclo^d			x				x				x				x		
Doxo^e					x				x				x				x
L-asp^f	x																
Pred^g	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x

^aCollette SA, Allstadt SD, Chon EM, *et al.* Treatment of feline intermediate- to high-grade lymphoma with a modified university of Wisconsin-Madison protocol: 119 cases (2004-2012). *Vet Comp Oncol* 2016;14:136-146.

^bNo chemotherapy administered on weeks 5, 10, 12, 14, 16, 18, 20, 22, and 24.

^cVincristine at 0.5 mg/m² IV.

^dCyclophosphamide at 227.3 to 242.4 mg/m² PO.

^eDoxorubicin at 0.89 to 1 mg/kg IV.

^fL-asparaginase at 10,000 U/m².

^gPrednisolone was continued throughout the entire 25-week protocol, starting with a dosage of 1.45 mg/kg, PO, SID tapered down to a chronic dosage of 0.42 mg/kg, PO, EOD for the patient's suspected asthma.